



Neutral composition of the Venus atmosphere measured by SPICAV/SOIR on board VENUS EXPRESS: A compendium

Ann Carine Vandaele, Arnaud Mahieux, Valérie Wilquet, Sarah Chamberlain, Bojan Ristic, Séverine Robert, Ian R. Thomas, Loïc Trompet, Jean-Loup Bertaux

► To cite this version:

Ann Carine Vandaele, Arnaud Mahieux, Valérie Wilquet, Sarah Chamberlain, Bojan Ristic, et al.. Neutral composition of the Venus atmosphere measured by SPICAV/SOIR on board VENUS EXPRESS: A compendium. International Venus Conference 2016, Apr 2016, Oxford, United Kingdom. insu-01297495

HAL Id: insu-01297495

<https://hal-insu.archives-ouvertes.fr/insu-01297495>

Submitted on 4 Apr 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

NEUTRAL COMPOSITION OF THE VENUS ATMOSPHERE MEASURED BY SPICAV/SOIR ON BOARD VENUS EXPRESS: A COMPENDIUM.

A. C. Vandaele, A. Mahieux, V. Wilquet, S. Chamberlain, B. Ristic, S. Robert, I. R. Thomas, L. Trompet, Royal Belgian Institute for Space Aeronomy, Brussels, Belgium (a-c.vandaele@aeronomie.be), J.-L. Bertaux, LATMOS, Paris, France.

Introduction: The Solar Occultation in the Infrared (SOIR) instrument on-board Venus Express is an infrared spectrometer covering the 2.2 - 4.3 μm spectral region. This instrument allows the detection of several key species of the Venus atmosphere, including CO_2 , CO, $\text{H}_2\text{O}/\text{HDO}$, HCl, HF and SO_2 . From the CO_2 density measurements, temperature is inferred giving information on the thermal structure of the atmosphere. Here we described the kind of data (profiles, latitudinal average, etc.) that have been obtained during the complete duration of the mission.

SOIR dataset: The Royal Belgian Institute for Space Aeronomy (BIRA-IASB) was in charge of SOIR's development and operations as well as its data pipeline.

BIRA-IASB carried out several studies on the composition of Venus mesosphere and lower thermosphere. The retrieval method has been described in detail in several papers introducing the solar occultation method [1], deriving CO_2 densities [2, 3] and trace gases such as CO [4, 5], HCl and HF [6], SO_2 [7-9], H_2O and HDO [10]. Temperature and total density, and therefore structure, has been retrieved from the CO_2 observations [2, 3, 11, 12]. Density and temperature profiles of the upper atmosphere of Venus (60 km to 170 km) at the terminator have been retrieved from SOIR's spectra using different assumptions, wherein the hydrostatic equilibrium and the local thermodynamical equilibrium in the radiative transfer calculations.

The SOIR spectra give also access to information on the aerosol content in the upper haze. Light extinction due to aerosols and cloud top altitudes have been obtained [13, 14] as well as the H_2SO_4 concentration in the droplets [15].

These results allow us to produce an Atmospheric model of Venus called Venus Atmosphere from SOIR measurements at the Terminator (VAST). Data obtained by SOIR will also contribute to update the Venus International Reference Atmosphere (VIRA) [16].

Recently, the treatment of the raw data to transmittance has been optimized, and a new dataset of spectra has been produced. All raw spectra (PSA level 2) as well as calibrated spectra (PSA level 3) have been delivered to ESA's Planetary Science Archive (PDS). Consequently the re-analysis of all spectra has been undergone. We will describe in detail the new derivation of the transmittances, and the updated dataset.

Results: In the following of the paper, we will describe in detail the data derived from the SOIR instrument, their coverage in terms of latitude, local solar time and altitude. As a general comment, solar occultation observations provide information on a very important region of the atmosphere, i.e. the terminator, the limit between the day and night sides of the planet. As a consequence the observations of SOIR correspond either to 6 am or 6 pm, even if at high latitude the definition of time starts to be meaningless. This region is crucial because it is a place of large longitudinal gradients in temperature and density.

Acknowledgments: The research program was supported by the Belgian Federal Science Policy Office and the European Space Agency (ESA – PRODEX program). The research was performed as part of the "Interuniversity Attraction Poles" programme financed by the Belgian government (Planet TOPERS). The authors also recognize the support from the FP7 EuroVenus project (G.A. 606798) as well as that of the HiResMIR International Scientific Coordination Network (GDRI). A. Mahieux would like to thank the FNRS (Belgium) for his Chargé de Recherches position.

References:

1. Vandaele, A.C., et al., *Composition of the Venus mesosphere measured by SOIR on board Venus Express*. J. Geophys. Res., 2008. **113**: p. doi:10.1029/2008JE003140.
2. Mahieux, A., et al., *Update of the Venus density and temperature profiles at high altitude measured by SOIR on board Venus Express*. Planet. Space Sci., 2015. **113-114**: p. 309-320.
3. Mahieux, A., et al., *Densities and temperatures in the Venus mesosphere and lower thermosphere retrieved from SOIR on board*

- Venus Express: Carbon dioxide measurements at the Venus terminator*. J. Geophys. Res., 2012. **117**(E07001): p. doi:10.1029/2012JE004058.
4. Vandaele, A.C., et al., *Carbon monoxide short term variability observed on Venus with SOIR/VEX*. Planet. Space Sci., 2015. **113-114**: p. 237-255.
 5. Vandaele, A.C., et al., *Carbon monoxide observed in Venus' atmosphere with SOIR/VEX*. Icarus, 2016. (**subm.**).
 6. Mahieux, A., et al., *Hydrogen Halides measurements in the Venus upper atmosphere retrieved from SOIR on board Venus Express*. Planet. Space Sci., 2015. **113-114**: p. 264-274.
 7. Belyaev, D., et al., *First observations of SO₂ above Venus' clouds by means of solar occultation in the infrared*. J. Geophys. Res., 2008. **113**: p. doi:10.1029/2008JE003143.
 8. Belyaev, D., et al., *Vertical profiling of SO₂ and SO above Venus' clouds by SPICAV/SOIR solar occultations*. Icarus, 2012. **217**(2): p. 740-751.
 9. Mahieux, A., et al., *Venus mesospheric sulfur dioxide measurement retrieved from SOIR on board Venus Express*. Planet. Space Sci., 2015. **113-114**: p. 193-204.
 10. Fedorova, A., et al., *HDO and H₂O vertical distributions and isotopic ratio in the Venus mesosphere by Solar Occultation at Infrared spectrometer onboard Venus Express*. J. Geophys. Res. . 2008. **113**(E00B22): p. doi:10.1029/2008JE003146.
 11. Mahieux, A., et al., *Densities and temperatures in the Venus mesosphere and lower thermosphere retrieved from SOIR onboard Venus Express: Retrieval technique*. J. Geophys. Res., 2010. **115**(E12014): p. 10.1029/2010JE003589.
 12. Mahieux, A., et al., *Rotational temperatures of Venus upper atmosphere as measured by SOIR on board Venus Express*. Planet. Space Sci., 2015. **113-114**: p. 347-358.
 13. Wilquet, V., et al., *Optical extinction due to aerosols in the upper haze of Venus: Four years of SOIR/VEX observations from 2006 to 2010*. Icarus, 2012. **217**(2): p. 875-881.
 14. Wilquet, V., et al., *Preliminary characterization of the upper haze by SPICAV/SOIR solar occultation in UV to mid-IR onboard Venus Express*. J. Geophys. Res., 2009. **114**: p. E00B42.
 15. Wilquet, V., et al. *SPICAV/SOIR mesospheric aerosols observations and characterization*. in *Geophysical Research Abstracts*, 16, 11491. 2014.
 16. Vandaele, A.C., et al., *Contribution from SOIR/VEX to the updated Venus International Reference Atmosphere (VIRA)*. Adv. Space Res., 2016. **57**: p. 443-458.